

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP013214

TITLE: Magnetic Field Induced Circular Photogalvanic Effect in InAs Quantum Wells

DISTRIBUTION: Approved for public release, distribution unlimited
Availability: Hard copy only.

This paper is part of the following report:

TITLE: Nanostructures: Physics and Technology International Symposium [9th], St. Petersburg, Russia, June 18-22, 2001 Proceedings

To order the complete compilation report, use: ADA408025

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP013147 thru ADP013308

UNCLASSIFIED

Magnetic field induced circular photogalvanic effect in InAs quantum wells

S. D. Ganichev^{†‡}, V. V. Bel'kov[‡], E. L. Ivchenko[‡], S. A. Tarasenko[‡],
 M. Sollinger[†], F.-P. Kalz[†], D. Weiss[†], J. Eroms[†] and W. Prettl[†]

[†] Institut für Experimentelle und Angewandte Physik,
 Universität Regensburg, D-93040 Regensburg, Germany
[‡] Ioffe Physico-Technical Institute, St Petersburg, Russia

We report on the first observation of a magnetic field induced circular photogalvanic effect (CPGE) in quantum wells (QWs). The experiments were carried out on (001)-MBE-grown *n*-InAs/AlGaSb QW structures with a 15 nm single InAs channel at 4.2 K. For optical excitation we used the $\lambda = 148 \mu\text{m}$ of a high power far-infrared pulsed NH_3 laser optically pumped by a TEA- CO_2 . The peak power of a single laser pulse was about 40 kW. The helicity P_{circ} of the incident light varied from -1 (left handed circular, σ_-) to +1 (right handed circular, σ_+) according to $P_{\text{circ}} = \sin 2\varphi$ where φ is the angle between the initial polarization plane and the optical axis of the $\lambda/4$ plate.

In the absence of a magnetic field, $\mathbf{B} = 0$, the irradiation of these semiconductor structures by far-infrared laser radiation results in a photocurrent, $\mathbf{j} \propto P_{\text{circ}}$, which reverses its sign by switching the helicity of radiation from left handed to right handed [1]. Due to the point-group symmetry C_{2v} of the studied QWs, the photogalvanic current at $\mathbf{B} = 0$ is only observed under oblique incidence. Here we demonstrate that the application of an external magnetic field, \mathbf{B} , in the interface plane induces a helicity-dependent current even at normal incidence. The current is proportional to B (up to 5 T) and inverts its direction with the reversal of the magnetic field. For the sake of brevity we refer to the effect under consideration as to the magneto-CPGE. For bulk materials this effect was theoretically treated in [2, 3] and observed in *p*-GaAs [4].

Phenomenologically, the magneto-CPGE is described by a third-rank tensor as

$$j_\alpha = \mu_{\alpha\beta\gamma} B_\beta i(\mathbf{E} \times \mathbf{E}^*)_\gamma = \mu_{\alpha\beta\gamma} E^2 B_\beta \hat{\mathbf{e}}_\gamma P_{\text{circ}}, \quad (1)$$

where \mathbf{E} is the amplitude of the electric field of the radiation, $E = |\mathbf{E}|$, and $\hat{\mathbf{e}}$ is a unit vector pointing in the direction radiation propagation.

In bulk crystals of the class T_d , the tensor $\mu_{\alpha\beta\gamma}$ has only one independent component $\mu \equiv \mu_{xyz}$, $\mu_{\alpha\beta\gamma} = \mu$ if $\alpha \neq \beta \neq \gamma$ and $\mu_{\alpha\beta\gamma} = 0$ otherwise. Hereafter we use the coordinate systems $x \parallel [100]$, $y \parallel [010]$, $z \parallel [001]$ and $x' \parallel [1\bar{1}0]$, $y' \parallel [110]$, $z \parallel [001]$. In a (001)-grown zinc-blende-lattice QW with non-equivalent normal and inverted interfaces, the point-group symmetry is reduced to C_{2v} . Under normal incidence of the light and for the magnetic field lying in the interface plane, the magneto-CPGE is described by two independent constants and, in the coordinate system (x', y', z) , can be presented as

$$\begin{aligned} \delta j_{x'} &= (\mu' + \mu) E^2 B_{x'} \hat{\mathbf{e}}_z P_{\text{circ}}, \\ \delta j_{y'} &= (\mu' - \mu) E^2 B_{y'} \hat{\mathbf{e}}_z P_{\text{circ}}. \end{aligned} \quad (2)$$

The photocurrent induced in the same geometry, $\hat{\mathbf{e}} \parallel z$, $\mathbf{B} \perp z$, in a bulk T_d -symmetry crystal or in a D_{2d} -symmetry QW with symmetrical interfaces is described by Eqs. (2)

assuming $\mu \neq 0$, $\mu' = 0$. In this case the directions of the vectors \mathbf{j} and \mathbf{B} are interconnected by the mirror reflection in the plane (110) if $\mu > 0$ or the plane (1 $\bar{1}$ 0) if $\mu < 0$. In particular, \mathbf{j} and \mathbf{B} are parallel (or antiparallel) when the magnetic field is applied along x' or y' and perpendicular when $\mathbf{B} \parallel x$ or $\mathbf{B} \parallel y$.

Another limiting case $\mu = 0$, $\mu' \neq 0$ is allowed not only by the C_{2v} symmetry but also by the polar uniaxial symmetry $C_{\infty v}$. The latter corresponds to the symmetry of a QW structure which is grown as if from isotropic compositional materials and has non-equivalent left- and right-hand-side interfaces. Note that if $\mu = 0$ then Eqs. (2) can be rewritten in the following two-dimensional vector form

$$\mathbf{j} = \mu' E^2 \mathbf{B} \hat{e}_z P_{\text{circ}}, \quad (3)$$

i.e. the vectors \mathbf{j} and \mathbf{B} are parallel irrespective to the in-plane orientation of \mathbf{B} .

The present experimental results are well described by Eq. (3) indicating that the symmetry of the investigated QW is $C_{\infty v}$. This is supported by the investigation of the circular photocurrent in the same structure under oblique incidence at $\mathbf{B} = 0$ for different geometries.

Acknowledgements

We thank for support the DFG, the RFBR and the NATO (Linkage Grant).

References

- [1] S. D. Ganichev, E. L. Ivchenko, S. N. Danilov, J. Eroms, W. Wegscheider, D. Weiss and W. Prettl, submitted to PRL.
- [2] E. L. Ivchenko and G. E. Pikus, *Problems in Modern Physics*, Nauka, Leningrad, 1980, p. 275.
- [3] E. L. Ivchenko, Yu. B. Lyanda-Geller and G. E. Pikus, *Sov. Phys. Solid State* **30**, 575 (1988).
- [4] A. V. Andrianov and I. D. Yaroshetskii, *Pis'ma Zh. Eksp. Teor. Fiz.* **40**, 131 (1984).